

7

# Healthy Eating for

6 to 24 month old children

Child Feeding  
Online Health Talk  
(In Cantonese only)



<https://s.fhs.gov.hk/u60sk>

Getting  
started

Introducing solid food



Department of Health

活出健康新方向  
For health



watch video



Website for the video:  
<http://s.fhs.gov.hk/oxfsd>

In the first few months of life, babies can only have breastmilk. For babies whose mothers are unable to breastfeed or whose parents have decided not to breastfeed, they can only have infant formula.

As their body systems and functions mature, most babies are ready to try solid foods when approaching 6 months of age.

This booklet helps you prepare for introducing solid foods to your babies.

## Contents

Introduce solid foods at around 6 months	3
Babies develop eating skills	4-5
How do you know that your baby is ready for eating solid foods ?	6
What are the suitable food choices for babies to start ?	7



How to prepare baby food	8
FAQs about first foods	9
Preparing for mealtime	10-11
How should parents feed babies with a spoon ?	12-13

How can we make mealtime enjoyable ?	14
Understand babies' signs of hunger and fullness	15-17
Foods and drinks that babies should avoid	18
Babies and food allergies	19





## Introduce solid foods at around 6 months

I am 6 months.  
I need eating other  
foods too.

### To satisfy babies' nutritional requirements

- After 6 months, babies have a higher demand for iron; breastmilk alone cannot satisfy their nutritional needs;
- Apart from breastmilk or infant formula, babies need a variety of foods to provide various nutrients to support their growth and development.



### To keep pace with babies' maturation and development

- Eating solid foods helps babies develop chewing;
- Giving babies a wide range of foods, food textures and tastes helps them accept family foods more easily;
- Babies are less likely to become picky eaters if they experience a variety of foods early.



To protect your baby from developing allergy,

**DO NOT feed him any solid foods before 4 months of age.**

What will happen when the introduction of solid food is delayed?

#### Risk of nutritional problems

Babies may not get adequate nutrients, e.g. iron and zinc, for growth, development and health.

#### Problems with eating habits

They may not accept a wide range of foods in the future and may thus become picky eaters. Some may refuse food with coarse textures.



## Babies develop eating skills

Babies still rely on breastmilk or formula milk as solid foods are first introduced. They gradually drink less milk while adapting to eat various nutritious foods. At around 2 years, most children dine with the family and eat family meals.



### How babies change in the way they eat

In the beginning



2 years

Sucking  
breastmilk or  
formula milk

Eating from  
a spoon and  
finger feeding  
themselves

Learning to  
use a cup

Feeding  
themselves with a  
spoon and eating  
with the family

### How babies accept new food textures

In the beginning



2 years

Smooth puree

Thicker,  
lumpier food




Soft, minced or  
chopped food

Food cut into  
small pieces



## Set a good example for babies

Babies tend to copy parents' eating behaviours and food choices. To help your baby form good eating habits, you should:


-  Consume a balanced diet and limit snacks of high sugar, fat and salt. Exercise regularly and maintain a healthy lifestyle;
-  Let your baby try foods that you like as well as foods that you dislike;
-  Avoid making negative comments about foods.





## Develop new skills

Babies are not only taking nutrients from foods but also learning **new skills**



### Learning to chew

-  Develops babies' abilities of chewing and swallowing.



### Learning to accept different tastes

-  Trying new foods allows babies to learn about the tastes of different foods;
-  Fosters babies' interest in foods.



### Developing the skill of feeding themselves

-  Babies learn to eat with spoon and drink from cup;
-  Babies learn to feed themselves.

### Eating as a family

-  Babies learn to eat properly at table;
-  Babies learn to eat socially.

### Developing relationships with family members

-  Good interaction during feeding enhances parent-child relationships;
-  Eating together allows you and your baby to enjoy a happy family time.



## How do you know that your baby is ready for eating solid foods?

🍎 Your baby can try solid foods if he has the following signs:

Watch Video



s.fhs.gov.hk/d1m2u

### Movements



- 🍌 Sitting well against the back of the chair;
- 🍌 Holding his head up;
- 🍌 Reaching out his hands to grasp objects.

### Eating behaviours



- 🥕 Showing interest in food;
- 🥕 Opening his mouth for a spoon;
- 🥕 Closing his lips over the spoon;
- 🥕 Able to swallow food.

- 🍎 The age when babies first show these signs highly varies, but most babies have these behaviours at about 6 months;
- 🍎 Premature babies can start eating solid foods when they are 6 months of age counting from the expected date of delivery and have these signs.

Discuss with your doctor or nurse if your baby does not show these signs when he is 7 months of age.

### Key points when starting to feed babies solid food

- 🍎 Babies still rely on milk feeding. Do not replace milk feedings at this stage;
- 🍎 Offer babies puree food 30 minutes before the usual feeding time;
- 🍎 To begin, give babies 1 - 2 teaspoonful puree at a time. Gradually increase the amount if babies eat well;
- 🍎 Choose foods that are iron-rich and can be made into smooth puree;
- 🍎 After trying smooth and runny food, babies can move on to try thicker food textures.



## What are the suitable food choices for babies to start?

First foods: Puree on spoon or finger foods? (text)

The first foods to start should be rich in iron and the texture should be smooth for babies to swallow. Choose foods that are soft and can be made into smooth puree.

Watch Video



s.fhs.gov.hk/5zj3e



s.fhs.gov.hk/qpr2k


### Grains

Infant cereal (rice, wheat, oat, etc.), strained congee




### Vegetables that can be easily pureed

Pumpkin, Chinese spinach, spinach, sweet potato



### Ripe and soft fruit




Banana, pear, peach, apple, papaya







### Meat, fish or eggs

Meat, egg yolk, fish, pork or chicken liver



-  As long as the foods are iron rich, there is no particular order that foods should be introduced;
-  You may offer iron-fortified infant cereal for 3 to 4 days and then add meat, vegetables or fruit puree. You can feed him the purees directly or mix them with infant cereal when serving.
-  Give vegetables for babies to try early so that they will accept other vegetables and fruits more easily.

### Iron rich first foods

-  Offer your baby foods rich in iron, such as egg, dark green leafy vegetables, livers, tofu, lentils, fish and meat. These foods can be pureed easily;
-  Take spoonful of these foods, such as egg yolk. Make into a puree or a paste with milk;
-  Or mix these foods with congee, infant cereal etc;
-  Please refer to the "7-day Healthy Meal Planning Guide for 6 to 24 month old children".

## How to prepare baby food

### Tools

Use a fine grater, strainer or sieve, or a blender to prepare puree for your baby.

### Baby food DIY

Let's learn about how to make food puree!



#### Suitable foods

Rice congee, Beans,  
Pork liver, Vegetable leaves

➡ Cooked and chopped finely

Vegetables like winter melon  
Carrot

➡ Cook until soft

Fruit

Egg yolk (hard boiled eggs)

#### Cooking methods

Grind the chopped pieces through a strainer with a spoon/rod

Ground into puree using a fine grater or press through a strainer

Scrape into puree using a spoon. For a finer texture, press it through a strainer

Mash cooked egg yolk with a fork. Add warm water to create a smooth texture

#### Food mash



### How to make infant cereal

1. Put infant cereal powder into bowl.
2. Mix well with warm water, breastmilk or formula milk, according to instructions on the package, to make a smooth puree.
3. Adjust the amount of water or milk to get a suitable texture.

Watch Video



s.fhs.gov.hk/z9yiz

For details, please refer to the "7-day Healthy Meal Planning Guide for 6 to 24 month old children".



## FAQs about first foods



### 1. How does infant cereal compare to rice congee?

- 🥦 Iron-fortified infant cereal contains more iron than rice congee (read the nutrition label on the package to check if iron is added);
- 🥦 Infant cereal is a convenient choice for some parents when babies first try solid food;
- 🥦 When babies get used to eating puree and demand more, parents can offer them congee with iron rich foods from the family food basket;
- 🥦 The iron provided by egg yolk, liver, green leafy vegetables, tofu, fish and meat can be absorbed easily. Changing tastes and textures of congee also help them adapt eating different foods and helps them chew.



### 2. Should mothers continue breastfeeding when solid food is introduced?

- 🥦 Mothers should continue breastfeeding. Breastmilk continues providing babies nutrients and protection against infection. This also benefits their long term health as well as the mothers'.

### 3. My baby start eating solid foods, should she continue taking vitamin D supplement?



- 🥦 Breastfed babies (exclusively breastfed or taking both breastmilk and formula milk) should continue to take vitamin D supplement until 12 months of age to ensure they have adequate vitamin D;
- 🥦 In addition, include a variety of vitamin D-containing foods such as fatty fish (e.g. salmon, sardine) and eggs in baby's diet. This can help further boost the intake of vitamin D.

### 4. My baby takes "No. 1" formula, does he have to change to "No. 2" formula?

- 🥦 He can continue with the "No. 1" infant formula he use to or change to "No. 2" formula after 6 months. Nevertheless, switching to "No. 2" formula is not necessary;
- 🥦 Babies can obtain sufficient nutrients from breastmilk or infant formula in addition to a diet including a variety of nutritious foods.




## Preparing for mealtime

Mealtime can be more enjoyable for you and your baby if some preparations are made.

### Four steps of preparing for mealtime

#### 1. "Warm-up" your baby

-  Before feeding, clean your baby's hands and face and set up a regular activity, say putting on the bib, chatting with him.



#### Purpose

To let babies know that "it is time for mealtime". This helps them to settle on a daily routine.

Clean my hands,  
put on a bib, sit me on  
my high chair. I know  
I am going to eat!






#### 2. Remove distractions

-  Turn off the TV, electronic devices and put away toys;
-  Keep your baby's attention on you and eating.

#### Purpose



To prevent problems resulting from eating while playing with toys or watching TV, such as:

-  Over-eating because of distractions;
-  Older babies may focus on watching TV and eat less;
-  Babies become less interested in feeding themselves.



Mum, my hands are clean.  
I'm ready to eat!

#### 3. Get yourself prepared

-  Your baby eats according to his needs. Do not worry on how much food he eats;
-  Prepare for the mess when your baby learns to self-feed. For example, cover the floor with newspaper.









Facing babies and responding to their needs can make them feel valued.

#### 4. The sitting position

-  Let your baby sit on a regular chair;
-  Sit with your baby face-to-face at the same level to encourage communication.

#### Purpose

Parents can observe babies more easily:



how he eats;



how he reacts to new foods;



his intention to self-feed;



whether he is full or hungry.

High chair



Avoid restraining your baby in the high chair for more than 1 hour at a time.

By getting used to sit on a regular location and seat for feeding, babies associate sitting on that chair and “it’s time for food”. It helps babies get prepared and settle themselves during mealtime.

The seat should be safe and comfortable. A high chair or a booster chair is ideal.

Booster chair



With a food tray attaching the chair, babies can reach food and feed themselves easily.

#### Parents should note the following if babies are fed in the following positions:

Sitting on your lap



It may be hard for you and your baby to see each other face-to-face.

On the flooring mat



Feeding may be difficult once he can crawl or walk. It is difficult for them to form the habit of "sitting down and eating".

On pushchair or baby walker




Babies can climb out of the pushchair, or move around with the baby walker. They will not learn to sit down to eat.

## How should parents feed babies with a spoon?

### Baby spoon

Wash baby's feeding utensils with soapy water or detergent. Sterilisation is not necessary.



### When should you feed your baby solid food?

- 🍎 Offer solids when your baby is relaxed. When he is too hungry or sleepy, he will be impatient to try solid foods;
- 🍎 Offer him a small amount of puree foods 30 minutes before the usual feeding time when he feels a bit hungry;
- 🍎 Let your baby taste the new food in the day time. It will be easier to observe any reactions.

### How to feed your baby food puree:



Get your baby prepared and bring him to the seat.  
(Please refer to p.10)

Let your baby see the food on the spoon.

When he opens his mouth, hold the spoon level on his lower lip.

When he closes his mouth, take the spoon out gently. Do not pour the food into his mouth.

### Hints :

1. If your baby cannot swallow the puree or he pushes out the spoon with his tongue, he is not ready for solid yet. Try again a week later;
2. Babies may drop some food from the side of their mouths at first. This will improve gradually as they become more skillful in chewing.





## How much to give

- 🍎 Begin with 1 - 2 teaspoons of puree once a day;
- 🍎 Offer more if your baby enjoys. Increase the amount and frequency gradually;
- 🍎 Offer your baby milk after feeding the solid foods. Do not cut milk feedings at this stage.

## When babies get used to eating the smooth puree

- 🍎 Increase the amount and the feeding of solid foods to 2 - 3 times a day;
- 🍎 Gradually adjust the food texture. Change to offer thick puree and soft mashed foods (please refer to the "Moving on" booklet);
- 🍎 Introduce new foods one at a time so as to build up babies' experience with food tastes. You can add new foods and serve with foods that baby has already tried.

Gradually adjust the food texture and increase the amount according to your baby's pace.



Egg yolk puree



Mashed egg yolk with milk

Add a new food one at a time



Tofu with sweet potato



Pork liver puree with rice cereal

## "Rules" for introducing new foods:

- 🍎 Food must be cooked thoroughly;
- 🍎 Introduce new foods one at a time;
- 🍎 Start with 1-2 teaspoons and increase the amount gradually. Try the food for 2 - 4 days before introducing another new food;
- 🍎 Observe for any allergic reactions (refer to P.19). If there is no allergic reaction, then try another new food.

## How can we make mealtime enjoyable?

### 1. Follow your baby's pace

- 🍉 Feed as fast or as slow as your baby eats;
- 🍉 When he loses interest in the food, call him softly to draw his attention;
- 🍉 Stop feeding when he shows signs of fullness.



### 2. Talk with your baby while feeding

- 🍉 Talk to him softly. Smile at him. These help him relax and eat better;
- 🍉 When your baby talks to you, respond to him promptly. This makes him happy;
- 🍉 Lack of communication makes him feel bored and anxious.

### 3. Allow babies to take part

- 🍉 When your baby shows interest in the spoon or food, let him touch and hold it or feed himself. Offer assistance if required.



### 4. Give babies encouragement

- 🍉 Praise your baby with words and body language when he tries new things;
- 🍉 Show him how to do something and he will be happier to try;
- 🍉 When he does well, praise him. Then, he knows that he has done the right thing.

## Understand babies' signs of hunger and fullness

- 🥑 Babies know how much they need to eat;
- 🥑 They show the signs of "I am hungry" or "I am full" through body language;
- 🥑 Most babies feel full within 15 to 30 minutes;
- 🥑 Let them lead the pace during feeding.



### Signs of hunger

- 1 Looking at food with interest;
- 2 Moving the head closer to the food and spoon;
- 3 Leaning towards the food;
- 4 Fussing or crying when too hungry.



### Signs of fullness

- 1 Less interested in eating;
- 2 Eating more and more slowly;
- 3 Turning her head away;
- 4 Pursing her mouth;
- 5 Spitting food out;
- 6 Pushing away or throwing the spoon and food;
- 7 Arching her back.



Babies know best how much they need to eat. No one knows better than your baby when his stomach is full.

### Hints:

If your baby gets full but you keep feeding him, your baby may:

- feel uncomfortable;
- link eating and discomfort together;
- struggle with you during mealtime, and he may end up eating less;
- over-eat which easily leads to obesity.





## Babies' appetite

Mum says that if I eat more, I shall grow faster and taller in the future! Is this true?



Babies require adequate nutrients to grow, but babies who eat more may not necessarily grow taller. Growth is also controlled by genes and affected by development during pregnancy. Overfeeding may lead to obesity and related health problems.

- 🍎 Actually, babies know how much to eat to get the right amount of nutrients for their growth and activities;
- 🍎 In the first three months of life, babies grow very fast and therefore they eat more;
- 🍎 When growth slows down, babies need less nutrients. Therefore, they eat less. It is normal that sometimes they do not seem to eat anything, but are still active and lively.

For a baby to grow healthily, the parents' job is to:

- Provide nutritionally appropriate and safe food;
- Feed babies according to their appetite.

My appetite varies from meal to meal !

- 🍎 Every baby is unique, so do not compare your baby's appetite with others' ;
- 🍎 Babies eat more after activities but sometimes refuse to eat when they get tired;
- 🍎 When they grow quickly, they eat more.



The ideal way to feed babies is to go according to their appetite. Do not insist that your baby eats the same amount at each meal.

## Common questions from parents



My baby is 4 months. His feeding is not regular, sometimes taking more, sometimes less. He gets distracted easily as well. If I feed him following his reactions, he might not get enough and cannot settle on the feeding schedule.

### Answer

It is very common for parents to have these worries.





You will be less anxious when you understand why your baby eats and behaves in this way.

### 1. Parents worry that babies may not get enough because they do not know the right amount to eat



My body has a well-built system telling me when and how much to eat. When I'm hungry, I'll tell you by my body language. If you feed me when I show you that I'm hungry, surely I shall get enough.





### 2. Parents worry babies may not get enough, as they are easily distracted

-  Parents should remove any objects that may distract children before mealtimes;
-  When baby is hungry, he usually eats faster and more attentively;
-  If a baby pauses and looks around, let him take a break. Then, call him to draw his attention to the food;
-  If he is still not interested in the food, that means he is full.

### 3. Parents worry that the amount of food taken changes from meal to meal

My appetite varies from meal to meal. When I ate less in the last meal, I shall eat more in the coming meal or earlier. If you feed me according to my full and hungry signs, I shall get all I need.

### 4. Parents worry that babies will not eat at regular times

-  In general, babies at 3 to 4 months have their own sleeping and eating patterns;
-  During daytime, most babies will not be hungry until 3 to 4 hours after feeding;
-  Feeding babies when they are not hungry disturbs their daily routine;
-  Forcing babies to follow your timetable may upset their regular eating patterns.



## Foods and drinks that babies should avoid

### 1. Drinks with added sugar will damage babies' teeth

#### Glucose water

Glucose drink does not help babies form the habit of water drinking.



#### Fruit Juice

- Infants do not need any fruit juice;
- They get more nutrients and dietary fibre from pureed or sliced fresh fruit;
- Toddlers 1 to 3 years of age should have **no more than 120ml fruit juice a day** if consumed.



#### Honey

- Honey may contain clostridium bacteria;
- Do not give honey to babies under 1 year old, as their immune systems are immature.



### 2. Tea, coffee & drinks containing caffeine, such as energy drinks or soft drinks

### 3. Fish with high levels of methylmercury

- This includes shark, swordfish, marlin, tuna (includes big eyes, bluefin, albacore, yellowfin species), king mackerel, splendid alfonsino, orange roughy, yellowback seabream and dash-and-dot goatfish.

### 4. Uncooked foods and unpasteurised milk products

### 5. Food that leads to choking easily

- Food with small and hard pieces: e.g. corn, peanuts, nuts, seeds;
- Food that is crispy or hard: e.g. candies, uncooked vegetables;
- Fish and meat with bones, and fruit with seeds.



### 6. Salt, soy sauce, chicken powder

- Babies may not easily accept bland-tasting food if they get used to salty tastes;
- Taking too much salt will increase their future risk for high blood pressure.





## Babies and food allergies

A food allergy is an immune response against certain food.

### Signs and symptoms of food allergies

Symptoms may occur within several hours or several days after eating the food.


- 🍎 Symptoms that occurs within several hours:  
Common symptoms:

- 🥕 Urticaria, worsening eczema;
- 🥕 Swollen eyes, tongue, face, mouth and lips;
- 🥕 Watery stools, vomiting.

Relatively rare but serious symptoms include:

- 🥕 Difficulty in breathing, fainting.

- 🍎 Some babies may have an allergic reaction 1 or 2 days later:  
e.g. eczema, persistent vomiting, wheezing, constipation, stomach pain.



Swollen eyes



Urticaria

© Department of Paediatrics and Adolescent Medicine, University of Hong Kong

### Foods that commonly cause food allergy

Milk products, eggs, peanuts, fish, shellfish, nuts, wheat (e.g. bread, biscuits), soy, bird's nest.



### When can babies try these foods?

- 🍌 Babies should not be given any solid foods before 4 months of age ;
- 🍌 Parents can introduce these foods when they start eating solid foods at about 6 months ;
- 🍌 Delaying or avoiding these foods cannot prevent babies from developing atopic dermatitis (eczema) or other allergic diseases ;
- 🍌 For babies with severe eczema or known food allergy, their parents should discuss with their paediatrician or family doctor before introducing these foods.

### If you think your baby has a food allergy:

- 🍎 Consult your family doctor as soon as possible. Stop giving the foods that you suspect is the problem until you have doctor's instruction;
- 🍎 If the reactions are serious, bring the baby to hospital immediately;
- 🍎 Babies diagnosed with food allergies must follow doctor's instruction on choosing food.

## Healthy eating for 6 to 24 month old children [Getting started] - Reminder to parents

- 🥕 At around 6 months, introduce solid foods to your baby to meet his nutrient requirement. Continue breastfeeding to allow him to obtain antibodies from the mother. This also benefits his long term health as well as the mother's;
- 🥕 Do not give your baby any food other than breastmilk or infant formula before 4 months of age. Consuming solid food too early, your baby will take less breastmilk and runs a higher risk of developing allergy;
- 🥕 When preparing his first foods, choose foods that are iron rich and from the family's food basket to provide variety (grains and cereals, vegetables, fruits, eggs, meat or fish);
- 🥕 Introduce new food one at a time and start with a small amount;
- 🥕 Your baby needs time to accept new foods. If he spits the food out or refuses to eat, do not force him, try again several days later;
- 🥕 Give your baby vitamin D supplement of 10 micrograms daily if he is breastfed.

If you have any queries about feeding your child,  
please contact your doctor and nurse.



To learn more about healthy eating for 6 to 24 month old children,  
please refer to the booklets "Moving On", "Ready to go"  
and "7-day Healthy Meal Planning Guide".

Booklet content will be updated from time to time. For the latest information,  
please refer to the website of the Family Health Service of the Department of Health.



**Family Health Service**

Website: [www.fhs.gov.hk](http://www.fhs.gov.hk)

24-hour information hotline: 2112 9900