

Pneumococcal Conjugate Vaccine

Pneumococcal infection

Pneumococcal infection represents a wide range of diseases caused by the bacterium *Streptococcus pneumoniae* (*S. pneumoniae* / pneumococcus). More than 90 serotypes have been identified. It can be transmitted by droplet spread and contact with respiratory secretions; direct contact is another way of transmission.

S. pneumoniae causes a wide spectrum of diseases, including:

- meningitis: it is a severe type of pneumococcal infection and usually presents with fever, stiff neck and mental confusion, causing long-term problems, such as hearing loss, or even death;
- pneumonia: it commonly presents with fever, shortness of breath, chills and productive cough, and may result in death in severe cases; and
- otitis media: it presents with fever, ear pain with or without ear discharge, and may lead to hearing loss in recurrent cases.

Pneumococcal Conjugate Vaccine (PCV)

A. Why get vaccinated?

Pneumococcal conjugate vaccine (PCV) can effectively protect against severe invasive infections due to the serotypes of *S. pneumoniae* contained in the vaccine. Young children are at risk of severe invasive pneumococcal infection (i.e. meningitis, bacteraemic pneumonia and septicaemia) and are advised to get vaccinated. In Hong Kong, PCV15 vaccine consisting of antigens against fifteen pneumococci serotypes) is included in the Hong Kong Childhood Immunisation Programme. More details about pneumococcal infection can be found in the Centre for Health Protection's website:

<https://www.chp.gov.hk/en/healthtopics/content/24/10584.html>



B. When should my child get vaccinated?

Children should receive 2 primary doses of PCV15 at 2 months and 4 months, followed by a booster dose of PCV15 at 12 months old.

C. The following individuals should NOT receive PCV

- Serious allergic reaction to previous dose of PCV
- Serious allergic reaction to diphtheria toxoid containing vaccine

D. What are the side effects?

- Most people have no serious reactions after receiving PCV.
- Occasionally there may be mild fever (usually occur within 3 days after vaccination) or slight redness or swelling around the injection site, but these will gradually subside in a few days. If fever or discomforts persist, or abnormal breathing is observed, please consult a doctor immediately.

E. How many types of PCV are available in HK and what are their protection?

There are more than 1 type of PCV available in the market of Hong Kong but no one PCV can cover all serotypes of *Streptococcus pneumoniae*. Personal and environmental hygiene are important measures in preventing pneumococcal infection.

If you have any query, please contact Maternal & Child Health Centre
of the Department of Health.